

Higher Condensation Theory

Liang Kong

SIQSE, Southern University of Science and Technology, China

Shenzhen-Nagoya Workshop, September 18–21, 2024

References: joint with Zhi-Hao Zhang, Jia-Heng Zhao, Hao Zheng, arXiv:2403.07813 (upcoming 2nd version!)

Anyon condensations in 2+1D


The mathematical theory of anyon condensation has a long history [Moore-Seiberg:1988-1989](#), [Bais-Slingerland:2002-2008](#), [Kapustin-Saulina:1008.0654](#), [Levin:1301.7355](#), [Barkeshli-Jian-Qi:1305.7203](#), ..., [Böckenhauer-Evans-Kawahigashi:math/9904109,0002154](#), [Kirillov-Ostrik:math/0101219](#), [Frölich-Fuchs-Runkel-Schweigert:math/0309465](#), [K.:1307.8244](#), ...

The mathematical theory of anyon condensation has a long history [Moore-Seiberg:1988-1989](#), [Bais-Slingerland:2002-2008](#), [Kapustin-Saulina:1008.0654](#), [Levin:1301.7355](#), [Barkeshli-Jian-Qi:1305.7203](#), ..., [Böckenhauer-Evans-Kawahigashi:math/9904109,0002154](#), [Kirillov-Ostrik:math/0101219](#), [Frölich-Fuchs-Runkel-Schweigert:math/0309465](#), [K.:1307.8244](#), ...

Theorem

Let \mathcal{C} and \mathcal{D} be the modular tensor categories (MTC) of anyons in two $2+1D$ topological orders. An anyon (or boson) condensation from \mathcal{C} to \mathcal{D} , which produces a gapped domain wall \mathcal{M} , is determined by a condensable E_2 -algebra A in \mathcal{C} .

- \mathcal{D} is the category of deconfined particles = the category of local A -modules (or E_2 - A -modules) in \mathcal{C} ; the trivial anyon $\mathbb{1}_{\mathcal{D}} \in \mathcal{D}$ is $A \in \mathcal{C}$;
 $\otimes_{\mathcal{D}} = \otimes_A$.
- \mathcal{M} is the category of (de)-confined particles = the category of right A -modules in \mathcal{C} .
- bulk-to-wall maps: $\mathcal{C} \xrightarrow{-\otimes A} \mathcal{M} \leftrightarrow \mathcal{D}$



Contrary to the physical intuitions that a phase transition between two phases are reversible, above mathematical description of an anyon condensation is not reversible because we have [Davydov-Müger-Nikshych-Ostrik:1009.2117](#).

$$\dim \mathcal{D} = \frac{\dim \mathcal{C}}{(\dim A)^2},$$

where $\dim A > 1$ for any non-trivial condensation.

Contrary to the physical intuitions that a phase transition between two phases are reversible, above mathematical description of an anyon condensation is not reversible because we have [Davydov-Müger-Nikshych-Ostrik:1009.2117](#).

$$\dim \mathcal{D} = \frac{\dim \mathcal{C}}{(\dim A)^2},$$

where $\dim A > 1$ for any non-trivial condensation.

It turns out that this phenomenon is a reflection of the fact that the 1-category of anyons (i.e., defects of codimension 2) does not include all topological defects in a 2+1D topological order [Kitaev-K.:1104.5047](#). In this talk, I will show that, by including all topological defects of codimension 1 and higher, we obtain a 2-category and a rather complete defect condensation theory, which is ready to be generalized to higher dimensions.

Category of topological defects

The mathematical theory of topological defects in an $n + 1$ D (potentially anomalous) topological order was developed in a series of works based on three guiding principles.

1. Remote Detectable Principle, [Levin:1301.7355](#), [K.-Wen:1405.5858](#)
2. Boundary-bulk relation, [Kitaev-K.:1104.5047](#), [K.-Wen-Zheng:1502.01690,1702.00673](#)
3. Condensation Completion Principle. [Carqueville-Runkel:1210.6363](#),
[Douglas-Reutter:1812.11933](#), [Gaiotto](#), [Johnson-Freyd:1905.09566](#), [Johnson-Freyd:2003.06663](#),
[K.-Lan-Wen-Zhang-Zheng:2003.08898](#)


Preceded by some earlier works [K.-Wen:1405.5858](#), [K.-Wen-Zheng:1502.01690](#), this theory was essentially established by Johnson-Freyd in 2020 [Johnson-Freyd:2003.06663](#) and was further developed and generalized to all quantum liquids in [K.-Zheng:2011.02859](#), [2107.03858](#), [2201.05726](#) .

We summarize the main results of this theory for an (potentially anomalous) $n+1$ D topological order \mathcal{C}^{n+1} .

1. The category of all topological defects (of codimension 1 and higher) form a fusion n -category \mathcal{C} (an \mathbb{E}_1 -fusion n -category). [Johnson-Freyd:2003.06663](#)
 - 1.1 0-morphisms (i.e., objects) in \mathcal{C} are 1-codimensional defects;
 - 1.2 1-morphisms in \mathcal{C} are 2-codimensional defects;
 - 1.3 k -morphisms are $(k + 1)$ -codimensional defects;
 - 1.4 n -morphisms are $(n + 1)$ -codimensional defects (i.e., instantons or 0D defects).
2. $\mathbb{1} \in \mathcal{C}$ labels the trivial 1-codimensional defect.
3. $\Omega\mathcal{C} := \text{hom}(\mathbb{1}, \mathbb{1})$ is the category of defects of codimension 2 and higher. It is braided fusion $(n - 1)$ -category or \mathbb{E}_2 -fusion $(n - 1)$ -category.
4. $\Omega^{k-1}\mathcal{C}$ is the category of defects of codimension k and higher. It is an \mathbb{E}_k -fusion $(n - k + 1)$ -category. [K.-Zheng:2011.02859](#)

I will explain the meaning of ' \mathbb{E}_1 ', ' \mathbb{E}_2 ' and ' \mathbb{E}_k '.

1. \mathcal{C} is E_1 -fusion: two 1-codimensional defects $x, y \in \mathcal{C}$ can be fusion in one direction $x \otimes^1 y$.
2. $\Omega\mathcal{C}$ is E_2 -fusion: two 2-codimensional defects $a, b \in \Omega\mathcal{C}$ can be fused in two orthogonal directions:



3. $\Omega^{k-1}\mathcal{C}$ is E_k -fusion: two k -codimensional defects $p, q \in \Omega^{k-1}\mathcal{C}$ can be fused in k orthogonal directions: $p \otimes^1 q, p \otimes^2 q, \dots, p \otimes^k q$.

Convention of notations:

- Topological orders: $A^{n+1}, B^{n+1}, C^{n+1}, \dots$;
- Category of all defects (codimension 1 or higher): $\mathcal{A}, \mathcal{B}, \mathcal{C}, \dots$;
- Category of defects of codimension 2 or higher: $\Omega\mathcal{A}, \Omega\mathcal{B}, \Omega\mathcal{C}, \dots$;
- Category of defects of codimension k or higher: $\Omega^{k-1}\mathcal{A}, \Omega^{k-1}\mathcal{B}, \Omega^{k-1}\mathcal{C}, \dots$.

Convention of notations:

- Topological orders: $A^{n+1}, B^{n+1}, C^{n+1}, \dots$;
- Category of all defects (codimension 1 or higher): $\mathcal{A}, \mathcal{B}, \mathcal{C}, \dots$;
- Category of defects of codimension 2 or higher: $\Omega\mathcal{A}, \Omega\mathcal{B}, \Omega\mathcal{C}, \dots$;
- Category of defects of codimension k or higher: $\Omega^{k-1}\mathcal{A}, \Omega^{k-1}\mathcal{B}, \Omega^{k-1}\mathcal{C}, \dots$.

If C^{n+1} is anomaly-free, then $\mathcal{C} = \Sigma\Omega\mathcal{C} = \text{Kar}(\text{B}\Omega\mathcal{C})$ ($n = 2$ [Carqueville-Runkel:1210.6363](#), [Douglas-Reutter:1812.11933](#); $n \geq 2$ [Gaiotto, Johnson-Freyd:1905.09566](#)).

$$\text{B}\Omega\mathcal{C} \simeq \begin{array}{c} \Omega\mathcal{C} = \text{hom}(\mathbb{1}, \mathbb{1}) \\ \curvearrowright \\ \mathbb{1} \end{array} \quad \Sigma\Omega\mathcal{C} = \text{Kar}(\text{B}\Omega\mathcal{C}) = \begin{array}{ccc} \Omega\mathcal{C} & & \Omega_x\mathcal{C} \\ \curvearrowright & \mathcal{M} & \curvearrowright \\ \mathbb{1} & \rightleftarrows & X \\ & \mathcal{M}^{\text{op}} & \end{array}$$

Convention of notations:

- Topological orders: $A^{n+1}, B^{n+1}, C^{n+1}, \dots$;
- Category of all defects (codimension 1 or higher): $\mathcal{A}, \mathcal{B}, \mathcal{C}, \dots$;
- Category of defects of codimension 2 or higher: $\Omega\mathcal{A}, \Omega\mathcal{B}, \Omega\mathcal{C}, \dots$;
- Category of defects of codimension k or higher: $\Omega^{k-1}\mathcal{A}, \Omega^{k-1}\mathcal{B}, \Omega^{k-1}\mathcal{C}, \dots$.

If C^{n+1} is anomaly-free, then $\mathcal{C} = \Sigma\Omega\mathcal{C} = \text{Kar}(\text{B}\Omega\mathcal{C})$ ($n = 2$ [Carqueville-Runkel:1210.6363](#), [Douglas-Reutter:1812.11933](#); $n \geq 2$ [Gaiotto, Johnson-Freyd:1905.09566](#)).

$$\text{B}\Omega\mathcal{C} \simeq \begin{array}{c} \Omega\mathcal{C} = \text{hom}(\mathbb{1}, \mathbb{1}) \\ \curvearrowright \\ \mathbb{1} \end{array} \quad \Sigma\Omega\mathcal{C} = \text{Kar}(\text{B}\Omega\mathcal{C}) = \begin{array}{ccc} \Omega\mathcal{C} & & \Omega_x\mathcal{C} \\ \curvearrowright & \mathcal{M} & \curvearrowright \\ \mathbb{1} & \rightleftarrows & X \\ & \mathcal{M}^{\text{op}} & \end{array}$$


Physically, $\mathcal{C} = \Sigma\Omega\mathcal{C}$ means that all 1-codimensional defects, which cannot be braided, can only be the condensation descendants of 2-codimensional defects, which can be braided and detectable by double braidings. [K.-Wen:1405.5858](#).

2+1D \mathbb{Z}_2 topological order TC^3 : We denote the fusion 2-category of topological defects in TC^3 by \mathcal{TC} . In this case, $\Omega\mathcal{TC}$ has four simple objects (or anyons) $1, e, m, f$:

$$e \otimes e \simeq m \otimes m \simeq f \otimes f \simeq 1, \quad f \simeq e \otimes m \simeq m \otimes e.$$

Mathematically, $\Omega\mathcal{TC}$ can be identified with the Drinfeld center $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ of $\text{Rep}(\mathbb{Z}_2)$. The fusion 2-category \mathcal{TC} consists of six simple objects (i.e. 1-codimensional topological defects) $1, \theta, ss, sr, rs, rr$. [K.-Zhang:2205.05565](#) θ is the invertible domain wall realizing the e - m duality.

\otimes	1	θ	ss	sr	rs	rr
1	1	θ	ss	sr	rs	rr
θ	θ	1	rs	rr	ss	sr
ss	ss	sr	$2ss$	$2sr$	ss	sr
sr	sr	ss	ss	sr	$2ss$	$2sr$
rs	rs	rr	$2rs$	$2rr$	rs	rr
rr	rr	rs	rs	rr	$2rs$	$2rr$



2+1D Ising topological order Is^3 : We denote the fusion 2-category of topological defects in Is^3 by \mathcal{J}_s . In this case, $\Omega\mathcal{J}_s = \{1, \psi, \sigma\}$ is the MTC of anyons.

It turns out that the only simple 1-codimensional topological defect in Is^3 is the trivial defect $\mathbb{1}$. [Fuchs-Runkel-Schweigert:hep-th/0204148](https://arxiv.org/abs/hep-th/0204148).

$$\mathcal{J}_s \simeq \Sigma\Omega\mathcal{J}_s \simeq B\Omega\mathcal{J}_s \simeq \begin{array}{c} \Omega\mathcal{J}_s \\ \curvearrowright \\ \mathbb{1} \end{array}$$

Theorem [Gaiotto, Johnson-Freyd:1905.09566](#): We denote the category of defects of the trivial topological order $\mathbf{1}^{n+1}$ by $n\text{Vec}$, which consists of only trivial k -dimensional defects $\mathbf{1}^k$ for $1 \leq k \leq n$ and their condensation descendants (i.e., defects that can be obtained from $\mathbf{1}^k$ via condensation).

(1) $\mathbf{1}^{0+1}$: $0\text{Vec} = \mathbb{C}$ (i.e., trivial fusion 0-category);


(2) $\mathbf{1}^{1+1}$: $1\text{Vec} := \Sigma\mathbb{C} = \text{Kar}(\text{BC}) = \{\mathbb{C}^{\oplus k}\}$ (i.e., trivial fusion 1-category);

$$\mathbb{1} = \mathbf{1}^1, \quad \begin{array}{c} \mathbb{C} \\ \curvearrowright \\ \mathbb{1} \end{array}$$


(3) $\mathbf{1}^{2+1}$: $2\text{Vec} = \Sigma\text{Vec} = \text{Kar}(\text{BVec}) = \{\text{Vec}^{\oplus k}\}$ (i.e., trivial fusion 2-category);

$$\mathbb{1} = \mathbf{1}^2, \quad \begin{array}{c} \text{Vec} \\ \curvearrowright \\ \mathbb{1} \end{array}$$

- (4) $\mathbf{1}^{n+1}$: $n\text{Vec} = \Sigma(n-1)\text{Vec} = \Sigma^n \mathbb{C} \neq \{(n-1)\text{Vec}^{\oplus k}\}$ for $n \geq 3$. For example, $3\text{Vec} = \{2\text{Vec}^{\oplus k}, \Sigma \mathcal{A} \mid \mathcal{A} \text{ is a multi-fusion 1-category}\} = \{\text{separable 2-categories}\}$.



A non-chiral topological order B^n
(i.e., admits a gapped boundary)
is a condensation descendant of $\mathbb{1} = \mathbf{1}^n$



Remark: The physical meaning of the $(n-1)$ -category \mathcal{M} is the category of gapped boundary conditions of B^n .

Definition (K.-Zheng:2011.02859)

It turns out that the functor $\text{hom}_{(n+1)\text{Vec}}(\bullet, -) : (n+1)\text{Vec} \rightarrow \text{Cat}_n^{\mathbb{C}}$ is fully faithful. We call an object (i.e., a \mathbb{C} -linear n -category) in the image of this functor a *separable n -category*. As a consequence, $(n+1)\text{Vec} = \text{Sep}_n$, where Sep_n denotes the category of separable n -categories.

$$\begin{array}{ccc}
 \begin{array}{c} (n-1)\text{Vec} \\ \curvearrowright \\ \mathbf{1}^n \\ \curvearrowleft \\ (n-1)\text{Vec} \end{array} & \begin{array}{c} \mathcal{M} \\ \xrightarrow{\quad} \\ \mathcal{M}^{\text{op}} \\ \xleftarrow{\quad} \end{array} & \begin{array}{c} \mathcal{B} \\ \curvearrowright \\ \mathbf{B}^n \\ \curvearrowleft \\ \mathcal{B} \end{array} \\
 & & \xrightarrow[\mathcal{A}=(n-1)\text{Vec}]{(n+1)\text{Vec} = \text{Sep}_n} \\
 & & \begin{array}{ccc}
 \text{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{A}) & & \text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{M}) \\
 \curvearrowright & \text{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{M}) & \curvearrowright \\
 \mathcal{A} & \xrightarrow{\quad} & \mathcal{M} \\
 \curvearrowleft & \text{Fun}_{\mathcal{A}}(\mathcal{M}, \mathcal{A}) & \curvearrowleft
 \end{array}
 \end{array}$$

Definition (K.-Zheng:2011.02859)

An E_m -multi-fusion n -category \mathcal{A} is a condensation-complete \mathbb{C} -linear E_m -monoidal n -category such that $\Sigma^m \mathcal{A}$ is a separable $(n+m)$ -category, i.e., $\Sigma^m \mathcal{A} \in (n+m+1)\text{Vec}$. If $\mathbb{1}_{\mathcal{A}}$ is simple, \mathcal{A} is an E_m -fusion n -category.

Condensations of topological defects

Theorem (K.-Zhang-Zheng-Zhao:2403.07813): Condensing a k -codimensional topological defect A in an $n+1$ D (potentially anomalous) topological order \mathcal{C}^{n+1} amounts to a k -step process.

- (1) The k -codimensional defect $A \in \Omega^{k-1}\mathcal{C}$ is condensable if it is equipped with the structure of a condensable E_k -algebra, i.e. an algebra equipped with compatible multiplications in k independent directions.

We first condense A along one of the transversal directions x^k , thus obtaining a $(k-1)$ -codimensional defect $\Sigma A := \text{RMod}_A(\Omega^{k-1}\mathcal{C}) \in \Sigma\Omega\mathcal{C}$.



- (2) It turns out that ΣA is naturally equipped with the structure of a condensable E_{k-1} -algebra, thus it can be further condensed along one of the remaining transversal direction x^{k-1} , thus obtaining a $(k-2)$ -codimensional defect $\Sigma^2 A := \text{RMod}_{\Sigma A}(\Omega^{k-2}\mathcal{C})$.



- (3) In the k -th step, condensing the 1-codimensional defect $\Sigma^{k-1} A$ along the only transversal direction defines a phase transition to a new $n+1$ D topological order D^{n+1} , which is Morita equivalent to C^{n+1} , and a gapped domain wall M^n .

$$Z(C)^{n+2} = Z(D)^{n+2}$$

$$C^{n+1} \quad M^n \quad D^{n+1}$$

$$\mathcal{D} \simeq \text{Mod}_{\Sigma^{k-1} A}^{E_1}(\mathcal{C}) = \text{BMod}_{\Sigma^{k-1} A | \Sigma^{k-1} A}(\mathcal{C})$$

$$(\mathcal{M}, m) = (\Sigma^k A := \text{RMod}_{\Sigma^{k-1} A}(\mathcal{C}), \Sigma^{k-1} A).$$

- (4) A k -codimensional deconfined topological defects in D^{n+1} form the category $\Omega^{k-1}\mathcal{D}$, which can be computed directly as the category of E_k - A -modules.

$$\Omega^{k-1}\mathcal{D} = \Omega^{k-1} \text{Mod}_{\Sigma^{k-1}A}^{E_1}(\mathcal{C}) \simeq \text{Mod}_A^{E_k}(\Omega^{k-1}\mathcal{C}),$$

A k -codimensional topological defect is deconfined iff it is equipped with a ' k -dimensional A -action', which defines the mathematical notion called an E_k -module over A or an E_k - A -module.

- (5) Similarly, the confined k -codimensional defects (confined to the wall M^n) can also be computed directly.

$$\Omega_m^{k-1}\mathcal{M} = \Omega^{k-1}\Sigma^k A = \Omega^{k-1} \text{RMod}_{\Sigma^{k-1}A}(\mathcal{C}) \simeq \text{RMod}_A(\Omega^{k-1}\mathcal{C}).$$

- (6) When C^{n+1} is anomaly-free (i.e., $\mathcal{C} = \Sigma\Omega\mathcal{C}$), the same phase transition, as a k -step process, can be alternatively defined by replacing the last two steps by a single step of condensing the E_2 -algebra $\Sigma^{k-2}A$ in the remaining two transversal directions directly.



The condensed phase D^{n+1} is determined by the category of E_2 -modules over $\Sigma^{k-2}A$.

$$\Omega\mathcal{D} \simeq \text{Mod}_{\Sigma^{k-2}A}^{E_2}(\Omega\mathcal{C}), \quad \mathcal{D} \simeq \Sigma\Omega\mathcal{D}, \quad \Omega_m\mathcal{M} = \text{RMod}_{\Sigma^{k-2}A}(\Omega\mathcal{C}).$$

When $n = 2$, this modified last step is precisely a usual anyon condensation in $2+1D$.

- (7) The condensable E_k -algebra A is called **Lagrangian** if $D^{n+1} = \mathbf{1}^{n+1}$.

Example: Consider 2+1D \mathbb{Z}_2 topological order TC^3 .

1. $A_e = 1 \oplus e$ is a Lagrangian E_2 -algebra in $\Omega\mathcal{TC}$. Condensing it along a line produces a string $\Sigma A_e = rr$, which can be further condensed to create the rough boundary of TC^3 .



2. $A_m = 1 \oplus m$ is a Lagrangian E_2 -algebra in $\Omega\mathcal{TC}$. Condensing it along a line produces a string $\Sigma A_m = ss$, which can be further condensed to create the smooth boundary of TC^3 .



Examples: $\mathcal{C}^{n+1} = \mathbf{1}^{n+1}$. In this case, the category of all defects $\mathcal{C} = n\text{Vec}$ (an E_1 -fusion n -category).

A condensable E_1 -algebra \mathcal{A} in $n\text{Vec}$ is precisely a multi-fusion $(n-1)$ -category. By condensing \mathcal{A} in $n\text{Vec}$, we obtain the condensed phase D^{n+1} and a gapped domain wall M^n .



D^{n+1} is a non-chiral topological order (i.e., admitting a gapped boundary) and a condensation descendant of $\mathbf{1}^{n+1}$

$$\mathcal{D} \simeq \text{BMod}_{\mathcal{A}|\mathcal{A}}(n\text{Vec}), \quad M^n = (\mathcal{M}, m) = (\text{RMod}_{\mathcal{A}}(n\text{Vec}), \mathcal{A})$$

$$\Omega\mathcal{D} = \text{Fun}_{\mathcal{A}|\mathcal{A}}(\mathcal{A}, \mathcal{A}) = \mathfrak{Z}_1(\mathcal{A}), \quad \Omega_m\mathcal{M} = \mathcal{A}$$

If $\mathfrak{Z}_1(\mathcal{A}) = (n-1)\text{Vec}$, then $\mathcal{D} = \Sigma\Omega\mathcal{D} = \Sigma(n-1)\text{Vec} = n\text{Vec}$ (i.e., $D^{n+1} = \mathbf{1}^{n+1}$). It means that a multi-fusion $(n-1)$ -category \mathcal{A} is **Lagrangian** iff $\mathfrak{Z}_1(\mathcal{A}) = (n-1)\text{Vec}$.

- 4 Since $\mathbf{C}^{n+1} = \mathbf{1}^{n+1}$ is anomaly-free, we can also condense a defect of codimension 2 directly. The category of 2-codimensional defects in $\mathbf{1}^{n+1}$ is $(n-1)\text{Vec}$. A condensable E_2 -algebra in $(n-1)\text{Vec}$ is precisely a braided fusion $(n-2)$ -category \mathcal{B} . By condensing \mathcal{B} directly along remaining two transversal directions, we obtain D^{n+1} and a gapped domain wall M^n .



D^{n+1} is a non-chiral topological order
(i.e., admitting a gapped boundary)
and a condensation descendant of $\mathbf{1}^{n+1}$

$$\Omega\mathcal{D} = \text{Mod}_{\mathcal{B}}^{E_2}((n-1)\text{Vec}), \quad \mathcal{D} = \Sigma\Omega\mathcal{D} \simeq \Sigma \text{Mod}_{\mathcal{B}}^{E_2}((n-1)\text{Vec}) \simeq \text{Mod}_{\Sigma\mathcal{B}}^{E_1}(n\text{Vec}),$$

$$\Omega_m\mathcal{M} = \text{RMod}_{\mathcal{B}}((n-1)\text{Vec}) = \Sigma\mathcal{B}, \quad \mathcal{M} = \Sigma\Omega_m\mathcal{M}.$$

- 4 Since $\mathbf{C}^{n+1} = \mathbf{1}^{n+1}$ is anomaly-free, we can also condense a defect of codimension 2 directly. The category of 2-codimensional defects in $\mathbf{1}^{n+1}$ is $(n-1)\text{Vec}$. A condensable E_2 -algebra in $(n-1)\text{Vec}$ is precisely a braided fusion $(n-2)$ -category \mathcal{B} . By condensing \mathcal{B} directly along remaining two transversal directions, we obtain D^{n+1} and a gapped domain wall M^n .

$$\Omega\mathcal{C} = (n-1)\text{Vec} \quad \left| \begin{array}{c} \Omega_m\mathcal{M} \\ \hline \Omega\mathcal{D} \end{array} \right.$$

D^{n+1} is a non-chiral topological order (i.e., admitting a gapped boundary) and a condensation descendant of $\mathbf{1}^{n+1}$

$$\begin{aligned} \Omega\mathcal{D} &= \text{Mod}_{\mathcal{B}}^{E_2}((n-1)\text{Vec}), & \mathcal{D} &= \Sigma\Omega\mathcal{D} \simeq \Sigma \text{Mod}_{\mathcal{B}}^{E_2}((n-1)\text{Vec}) \simeq \text{Mod}_{\Sigma\mathcal{B}}^{E_1}(n\text{Vec}), \\ \Omega_m\mathcal{M} &= \text{RMod}_{\mathcal{B}}((n-1)\text{Vec}) = \Sigma\mathcal{B}, & \mathcal{M} &= \Sigma\Omega_m\mathcal{M}. \end{aligned}$$

When $n = 3$, every fusion 2-category is Morita equivalent to $\Sigma\mathcal{A}$ for a braided fusion 1-category \mathcal{A} . [Décoppet:2208.08722](#) It means that all non-chiral 3+1D topological orders can be obtained from $\mathbf{1}^4$ by condensing defects of codimension 2 (i.e., strings).

When $\mathcal{C}^{n+1} = \mathcal{D}^{n+1} = \mathbf{1}^{3+1}$, the category of strings is $\Omega\mathcal{C} = 2\text{Vec}$. The gapped domain wall M^3 is precisely a 2+1D anomaly-free topological order, which can be described by a pair (\mathcal{B}, c) for a modular tensor category \mathcal{B} .



By rolling up the 2+1D anomaly-free topological order M^3 , we obtain a string-like defect in $\mathbf{1}^4$, which is precisely the object $\mathcal{B} \in \Omega\mathcal{C} = 2\text{Vec}$. By condensing this string, we obtain $\Omega\mathcal{D} = \text{Mod}_{\mathcal{B}}^{\mathbb{E}_2}(2\text{Vec}) \simeq \Omega \text{Mod}_{\Sigma\mathcal{B}}^{\mathbb{E}_1}(3\text{Vec}) \simeq \Omega 3\text{Vec} = 2\text{Vec}$. Therefore, $\mathcal{D}^4 = \mathbf{1}^4$ and \mathcal{B} is a Lagrangian \mathbb{E}_2 -algebra in 2Vec .

(3). \mathcal{C}^{n+1} is anomaly-free, i.e. $\mathfrak{Z}_1(\mathcal{C}) = n\text{Vec}$. In this case, we have

$$\mathcal{C} = \Sigma\Omega\mathcal{C} = \text{RMod}_{\Omega\mathcal{C}}(n\text{Vec})$$

provides a concrete coordinate system to the non-degenerate fusion n -category \mathcal{C} .

Theorem ([Brochier-Jordan-Synder:1804.07538](#), [K.-Zhang-Zhao-Zheng:2403.07813](#))

*An indecomposable condensable E_1 -algebra in \mathcal{C} are precisely an indecomposable multi-fusion $(n-1)$ -category \mathcal{A} equipped with a central functor $L : \Omega\mathcal{C} \rightarrow \mathcal{A}$ (i.e., a braided monoidal functor $\phi : \Omega\mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{A})$). When $\phi : \Omega\mathcal{C} \rightarrow \mathfrak{Z}_1(\mathcal{A})$ is a braided equivalence, the condensable E_1 -algebra \mathcal{A} is **Lagrangian** in the sense that $D^{n+1} = \mathbf{1}^{n+1}$.*


Condensing \mathcal{A} in \mathcal{C} produces a condensed phase D^{n+1} and a gapped domain wall M^n :

$$\mathcal{D} \simeq \text{Mod}_{\mathcal{A}}^{E_1}(\mathcal{C}), \quad \mathcal{M} \simeq \text{RMod}_{\mathcal{A}}(\mathcal{C}) \simeq \text{RMod}_{\mathcal{A}}(n\text{Vec}), \quad m = \mathcal{A},$$

Question: Does the abstract data of $L : \Omega\mathcal{C} \rightarrow \mathcal{A}$, which defines an abstract algebra, has a direct physical meaning?

Physical Meaning of this Theorem:

$$\begin{aligned}\mathcal{M} &\simeq \text{RMod}_{\mathcal{A}}(\mathcal{C}) \simeq \text{RMod}_{\mathcal{A}}(n\text{Vec}), \\ m &= \mathcal{A}, \\ \Omega_m \mathcal{M} &= \text{RMod}_{\mathcal{A}}(\mathcal{A}, \mathcal{A}) \simeq \mathcal{A}.\end{aligned}$$




Take home message: The multi-fusion higher category $\Omega_m \mathcal{M}$ (of all defects on the domain wall M^n) is a condensable E_1 -algebra in \mathcal{C} .

Example: $C^3 = TC^3$: The topological defects in TC^3 form the fusion 2-category

$$\mathcal{TC} \simeq \Sigma(\Omega\mathcal{TC}) = \text{RMod}_{\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))}(2\text{Vec}).$$


A condensable E_1 -algebra in \mathcal{TC} are precisely a mult-fusion categories \mathcal{A} equipped with a central functor $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) \rightarrow \mathcal{A}$.

1. $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) \xrightarrow{\text{id}} \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$ defines a condensable E_1 -algebra $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))$. It is just the trivial condensable E_1 -algebra, i.e., tensor unit $\mathbb{1}$ of \mathcal{TC} .
2. $f : \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) \rightarrow \text{Rep}(\mathbb{Z}_2)$ defines a condensable E_1 -algebra $\text{Rep}(\mathbb{Z}_2)^{\text{op}} = \text{Rep}(\mathbb{Z}_2)$ in \mathcal{TC} , which is precisely the 1-codimensional defect ss in \mathcal{TC} .

$$\text{Mod}_{\text{Rep}(\mathbb{Z}_2)}^{E_1}(\mathcal{TC}) \simeq 2\text{Vec},$$


Since the condensed phase is trivial, ss is a Lagrangian condensable E_1 -algebra in \mathcal{TC} .

- 3 $\mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2)) \xrightarrow{\cong} \mathfrak{Z}_1(\text{Vec}_{\mathbb{Z}_2}) \rightarrow \text{Vec}_{\mathbb{Z}_2}$ defines a condensable E_1 -algebra $\text{Vec}_{\mathbb{Z}_2}^{\text{op}} = \text{Vec}_{\mathbb{Z}_2}$, which is precisely the 1-codimensional defect rr in \mathcal{TC} . We have

$$\text{Mod}_{\text{Vec}_{\mathbb{Z}_2}}^{E_1}(\Sigma \mathfrak{Z}_1(\text{Rep}(\mathbb{Z}_2))) \simeq 2\text{Vec},$$


Since the condensed phase is trivial, rr is a Lagrangian condensable E_1 -algebra in \mathcal{TC} .

- 4 Consider double Ising 2+1D topological order ($\Omega\mathcal{I}s$ is the Ising MTC).

$$A := \mathbb{1} \boxtimes \mathbb{1} \oplus \psi \boxtimes \psi \in \Omega\mathcal{I}s \boxtimes \Omega\mathcal{I}s^{\text{op}} \simeq \mathfrak{Z}_1(\Omega\mathcal{I}s) \quad (1)$$

has a condensable E_2 -algebra in $\mathfrak{Z}_1(\Omega\mathcal{I}s)$. Then the central functor

$$\{1, e, m, f\} = \Omega\mathcal{TC} = \text{Mod}_A^{E_2}(\mathfrak{Z}_1(\Omega\mathcal{I}s)) \hookrightarrow \text{RMod}_A(\mathfrak{Z}_1(\Omega\mathcal{I}s))^{\text{op}} = \{1, e, m, f, \chi_{\pm}\} = \mathcal{K}$$

[Chen-Jian-K.-You-Zheng:1903.12334](#) defines a condensable E_1 -algebra in \mathcal{TC} . Condensing it in \mathcal{TC} produces the 2+1D double Ising topological order as the condensed phase.

$$\text{Mod}_{\mathcal{K}}^{E_1}(\mathcal{TC}) \simeq \mathcal{I}s \boxtimes \mathcal{I}s^{\text{op}}.$$

A higher dimensional example: Consider the $n+2$ D G -gauge theory GT_G^{n+2} . The category of 2-codimensional defects in GT_G^{n+2} was conjectured in [K.-Tian-Zhou:1905.04644](#) to be:

$$\Omega \mathcal{GT}_G^{n+2} = \mathfrak{Z}_1(n\text{Rep}(G)) \simeq \mathfrak{Z}_1(n\text{Vec}_G) \simeq \bigoplus_{[h] \in \text{Cl}} n\text{Rep}(C_G(h))$$

The fusion $(n+1)$ -category \mathcal{GT}_G^{n+2} has two coordinate systems:

$$\mathcal{GT}_G^{n+2} = \text{RMod}_{\mathfrak{Z}_1(n\text{Rep}(G))}((n+1)\text{Vec}), \quad \mathcal{GT}_G^{n+2} = \text{BMod}_{n\text{Rep}(G)|n\text{Rep}(G)}((n+1)\text{Vec})^{\text{op}},$$

We have $\Omega \mathcal{GT}_G^{n+2} = \mathfrak{Z}_1(n\text{Rep}(G))$ and

$$\Omega^k \mathcal{GT}_G^{n+2} = (n-k+1)\text{Rep}(G) \quad \text{for } k \geq 2$$

When $k = n$, $\Omega^n \mathcal{GT}_G^{n+2} = \text{Rep}(G)$ is the 1-category of particles.

Let $H < G$ be a subgroup of G . The composite particle

$$A = \text{Fun}(G/H) \quad (\text{i.e., } \mathbb{C}\text{-valued functions on } G/H)$$

is an E_{n+1} -algebra in $\Omega^n \mathcal{G}\mathcal{T}_G^{n+2} = \text{Rep}(G)$. By condensing the A -particles, we mean the following procedures.

(1) We first condensing the A -particle along a line, we obtain a string

$$\Sigma A = \text{RMod}_A(\text{Rep}(G)) \simeq \text{Rep}(H) \in 2\text{Rep}(G) = \Omega^{n-1} \mathcal{G}\mathcal{T}_G^{n+2}.$$

(2) We further condense the ΣA -string along one of the remaining transversal directions, we obtain a membrane:

$$\Sigma^2 A = \text{RMod}_{\Sigma A}(2\text{Rep}(G)) \simeq \text{RMod}_{\Sigma A}(2\text{Vec}) \simeq 2\text{Rep}(H) \in 3\text{Rep}(G) = \Omega^{n-2} \mathcal{G}\mathcal{T}_G^{n+2}.$$

(3)

$$\begin{aligned}\Sigma^{n-1}A &= (n-1)\mathrm{Rep}(H) \in n\mathrm{Rep}(G) \hookrightarrow \mathfrak{Z}_1(n\mathrm{Rep}(G)). \\ \Sigma^n A &= \mathrm{RMod}_{\Sigma^{n-1}A}(\mathfrak{Z}_1(n\mathrm{Rep}(G))) \in \Sigma\mathfrak{Z}_1(n\mathrm{Rep}(G)).\end{aligned}$$

Translate $\Sigma^n A$ into an E_1 -algebra in the second coordinate system of \mathcal{GT}_G^{n+2} . It is defined by the following monoidal functor:

$$\begin{aligned}n\mathrm{Rep}(G) &\rightarrow n\mathrm{Rep}(G) \boxtimes_{\mathfrak{Z}_1(n\mathrm{Rep}(G))} \mathrm{RMod}_{\Sigma^{n-1}A}(\mathfrak{Z}_1(n\mathrm{Rep}(G))) \\ &\simeq \mathrm{RMod}_{\Sigma^{n-1}A}(n\mathrm{Rep}(G)) \simeq n\mathrm{Rep}(H).\end{aligned}$$

(4) By condensing $\Sigma^n A = n\mathrm{Rep}(H)$, we obtain $D^{n+2} = \mathrm{GT}_H^{n+2}$ as the condensed phase:

$$\mathcal{D} \simeq \mathrm{Mod}_{n\mathrm{Rep}(H)^{\mathrm{op}}}^{E_1}(\mathcal{GT}_G^{n+2}) \simeq \mathrm{Mod}_{n\mathrm{Rep}(H)^{\mathrm{op}}}^{E_1}((n+1)\mathrm{Vec}) \simeq \mathcal{GT}_H^{n+2}.$$

As a consequence, we provides the precise mathematical theory behind the folklore that breaking the G -gauge “symmetry” in GT_G^{n+2} to a subgroup H gives the H -gauge theory GT_H^{n+2} .

Conclusion and outlooks

1. One can see that the theory of defect condensation is precisely a mathematical theory of higher representations of higher algebras or higher Morita theories. There will be a mathematical companion of this paper, in which we develop a mathematical theory of condensable E_k -algebras.
2. It is possible to developed a mathematical theory of condensations of gapless but liquid-like defects based on the theory of gapped/gapless quantum liquids [K.-Zheng:1705.01087,1905.04924,1912.01760,2011.02859](#), which is a prehistorical theory of SymTO/SymTFT based on the so-called 'topological Wick rotation'.
3. Although a new paradigm is emerging, as far as I can tell, it is still far from being complete. A complete paradigm demands an entirely new calculus, in which we are still in the beginning stage to understand integers. It means that there are a lot of exciting problems to work on in the coming future.

Thank you!